

# Cholesterol Oxidase “Amano” 6E

(Cholesterol: oxygen oxidoreductase, EC 1.1.3.6)

Cholesterol Oxidase “Amano” 6E is a cholesterol oxidase preparation extracted from bacteria.

## Catalysis



## Specification and Preparation

Activity:	Cholesterol oxidase activity	$\geq$ 10 u/mg (Amano new method)
Contaminant:	Catalase activity	$\leq$ 0.5%
Appearance:	Light yellow to yellowish orange powder, lyophilized	
Additives:	Saccharose	

## Characteristics

1. Molecular weight: 64,000 (SDS-PAGE)
2. Isoelectric point: 4.7
3. Km:  $3.0 \times 10^{-5}$  M
4. Optimum pH: 7.0
5. pH stability: 5.0-9.0 (37°C, 1hr)
6. Optimum temperature: 45-50°C
7. Thermal stability: up to 40°C (4hr, dissolved in purified water)
8. Activator: Sodium cholate
9. Stabilizer: Sodium cholate
10. Application: Used for enzymatic determination of cholesterol in serum by coupling with cholesterol esterase in clinical diagnosis.

## Expiration (Storage)

24 months from the date of analysis when stored at 5°C or below in a dry place under sealed conditions.

The information and recommendations contained herein are to the best of our knowledge reliable according to the current scientific and technical level. However, depending upon use method and/or condition, nothing herein is to be construed as a warranty or representation in respect otherwise, including freedom from patent infringement. Users shall make their own test and investigation for their particular purpose. We do not accept any liability for any loss, damage or infringement arising from the use of information and recommendations contained herein.


## Safe Handling

1. Do not inhale.
2. In case of direct contact with skin or eyes, immediately wash or rinse with plenty of water.
3. Please refer to SDS for more details.


## General properties

The following results demonstrate the activities of enzyme solution prepared in various buffers. Enzyme activity may vary under different experimental conditions.


### pH and Activity




### pH stability



### Temperature and Activity




### Thermostability



### Substrate Specificity


Substrate	Relative Activity (%)
Cholesterol	100
Dehydroisoandrosterone	13
Pregnenolone	32
β-Sitosterol	82
Stigmasterol	64
Estradiol	3
Vitamin D3	0
Cholic acid	0
Androsterone	0
Cholesterol linoleate	0
Lanosterol	0

### Stability (powder form)



# Assay method of Cholesterol oxidase activity

## Principle



The appearance of quinoneimine dye formed by coupling with 4-aminoantipyrine and phenol is measured at 500 nm by spectrophotometry.

## Unit Definition

One unit is defined as the enzyme quantity which oxidizes one  $\mu$  mole of cholesterol per minute under the conditions described below

## Reagents

- A. 4-Aminoantipyrine solution (5 mg/ml deionized water)
- B. Triton X-100 stock solution (50 mg/ml deionized water)
- C. Phenol solution (50 mg/ml deionized water)
- D. 0.1M Phosphate buffer ( $\text{KH}_2\text{PO}_4$ -NaOH, pH 7.0)
- E. 0.1% Triton-X Solution  
Dilute Triton X-100 stock solution (B) to 0.1% concentration with deionized water.
- F. Aminoantipyrine-phenol solution  
Mix 5.5 ml of 4-Aminoantipyrine solution (A), 5.5 ml of Phenol solution (C) and 29 ml of TritonX-100 stock solution (B), then fill up to 500 ml with 0.1 M Phosphate buffer (D). After preparation, release the lid of the container and allow the solution to stand under refrigeration for 24 hours before use. (Can be used for 3 months in light-shielded and refrigerated storage)
- G. Substrate solution  
Weigh 23 mg of cholesterol and dissolve in 10 ml of 2-propanol. (Can be used for 5 days at room temperature)
- H. Peroxidase solution  
Dissolve 330 units of Peroxidase (Amano Enzyme) in 2ml of chilled 0.1 M Phosphate buffer (D).  
(Can be used for 2 weeks at 2-8°C)
- I. Working solution  
Mix 1ml of Substrate solution (G) and 58 ml of Aminoantipyrine-phenol solution (F), then add 1 ml of Peroxidase solution (H) and mix.  
(Can be used for 18 days in light-shielded and refrigerated storage)
- J. Enzyme solution  
Weigh 20 mg of Cholesterol Oxidase "Amano" 6E and dissolve in 5 ml of deionized water and keep the solution at  $37 \pm 0.5^\circ\text{C}$  in a water bath for 4 hours. Then, after cooling down in an ice bath, dilute the solution with chilled 0.1% Triton X-100 solution (E). Enzyme solution should be prepared so that the value of  $\Delta\text{OD}/\text{minute}$  becomes in the range of  $0.020 \pm 0.005$ .

## Procedure

Put 3ml of Working solution (I) into a quartz cell (d = 10 mm) and keep at 37±0.5°C for 10 minutes. Then, pipette 0.1 ml of Enzyme Solution (J) into the quartz cell and mix well immediately. Keep the reaction mixture at 37±0.5°C. Exactly at 3 minutes and 6 minutes after the addition of Enzyme solution (J), measure the absorbances of the reaction mixture at 500 nm (A3 and A6). As a blank, pipette 0.1% Triton X-100 solution (E) into another quartz cell instead of Enzyme solution (J) and take the same procedure described above (Ab3 and Ab6).

## Calculation

$$\text{Cholesterol Oxidase activity (u/mg)} = \frac{(A6 - A3) - (Ab6 - Ab3)}{3} \times \frac{1}{13.6} \times 2 \times 3.1 \times \frac{n}{0.1}$$

- 3: Reaction time
- 13.6: Millimolar absorption coefficient of quinoneimine dye
- 2: Conversion factor (1 mole of quinoneimine dye equivalent to 2 mole of cholesterol)
- 3.1: Volume of the reaction mixture
- 0.1: Volume of Enzyme solution
- n: Dilution factor of Enzyme solution

## Contact

<i>Area</i>	<i>Branch</i>	<i>Location</i>	<i>E-mail</i>
North, Central, South America	Amano Enzyme U.S.A. Co., Ltd.	Illinois, U.S.A.	aeu.sales@amano-enzyme.com
Europe, the Middle East and Africa	Amano Enzyme Europe Limited	Oxfordshire, U.K.	aee.sales@amano-enzyme.com
Asia Pacific	Amano Enzyme Asia Pacific Co., Ltd.	Pathum Thani, THAILAND	aeap.sales@amano-enzyme.com
China	Amano Enzyme Manufacturing (China), Ltd. Shanghai Branch	Shanghai, P.R.CHINA	shanghai@amano-enzyme.com.cn
Japan, Headquarters	Amano Enzyme Inc.	Nagoya, JAPAN	